

EXPERIMENT EXAMINATION RULES

1. You are NOT allowed to bring any personal items into the examination room, except for the water bottle, personal medicine or approved personal medical equipment.
2. You must sit at your designated desk and must always ensure that the camera/supervisor is able to clearly see you.
3. Check the stationery items (pen, calculator, and scrap paper) provided by the organizers.
4. Do NOT start answering the questions before the “START” signal.
5. You are NOT allowed to leave the examination room during the examination except in an emergency in which case you will be accompanied by a supervisor.
6. If you need to visit the bathroom, please raise your hand.
7. Do NOT disturb other competitors. If you need any assistance, raise your hand and wait for a supervisor to come.
8. You can ONLY ask questions and discuss the experiments with your own team members. You must STAY at your table until the end of the time allocated for the experiments, even if you have finished the experiments or do not wish to continue.
9. At the end of the examination time you will hear the “STOP” signal. Do NOT write anything more on the answer sheet after this stop signal.
10. Arrange the exam, answer sheets, and the stationary items (pen, calculator, and scrap paper) neatly on your desk. All the pages will be submitted back including the question paper and rough work.
11. Do not leave the room before all the answer sheets have been collected and Supervisors give you a signal to leave the room

Experiment



G0-2

TWN Fu (Chinese Taipei)

EXPERIMENT EXAMINATION INSTRUCTIONS

1. After the “START” whistle, you will have 3 hours to complete the exam.
2. Please confirm that the student code printed on your question paper and answer sheet is same as your student code provided to you by organizers.
3. You have 18 pages of answer sheets—including the front page. Raise your hand, if you find any sheets missing, after the “START” signal is given.
4. Read the experimental procedures and questions carefully and write your answers in the corresponding boxes of the answer sheets.
5. When units are provided in the answer sheets, you have to write the answers correctly for the units.
6. Always show your calculations if room for this is provided. If you do not show your calculations, no points are awarded for the question.
7. You MUST wear a Lab Coat and Safety Glasses during the experiments.
8. You will be provided with two sets of answer sheets. Only one on which "Final answer sheets" is written will be evaluated.
9. You can use the "Draft answer sheet" for noting down your initial readings and scrap work, but they will NOT be evaluated.
10. The number of points that can be obtained is indicated for each question.
11. The total number of questions is 5. Check if you have a complete set of the test questions sheets(13pages, page 6–page 18)after the “START” signal is given. Raise your hand, if you find any sheets missing.
12. Useful information for answering the questions (atomic masses, constants, and formulas) is provided on page 4.
13. Always show your calculations. If you do not show your calculations, no points are awarded for the question.
14. You should write your final answers down in the appropriate number of significant digits.

Experiment



G0-3

TWN Fu (Chinese Taipei)

List of Constants

- Avogadro' s constant $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$
- Electronic charge $e = 1.602 \times 10^{-19} C$
- Molar gas constant $R = 8.314 \frac{J}{\text{mol } K}$
- Molar gas constant $R = 0.082 \frac{L \text{ atm}}{\text{mol } K}$
- Molar gas constant $R = 1.982 \frac{cal}{\text{mol } K}$
- Planck' s constant $h = 6.626 \times 10^{-34} Js$
- Speed of light (in vacuum) $c = 2.998 \times 10^8 \frac{m}{s}$
- 1 atomic mass unit $1u = 931.5 \frac{MeV}{c^2}$
- Specific heat of water $= 4.2 \frac{kJ}{kg \cdot K}$
- 1 Dalton $= 1.661 \times 10^{-27} kg$
- $1 eV = 1.602 \times 10^{-19} J$
- $1 \text{ bar} = 10^5 Pa = 10^5 Nm^{-2} = 1 \text{ atm}$
- Acceleration due to gravity $g = 9.8 \frac{m}{s^2}$
- 1 Litre $= 10^3 cm^3 = 10^{-3} m^3$
- Density of water $= 1000 \frac{kg}{m^3}$
- Mechanical equivalent of heat $= 4.186 \frac{J}{Cal}$
- 1 atmosphere $= 10^5 Pascal$
- Refractive index of water $= 1.33$
- 1 mL $= 1 cm^3$

Experiment

Group																			
I	II											III	IV	V	VI	VII	VIII		
Key atomic number atomic symbol name relative atomic mass												1 H hydrogen 1							2 He helium 4
3 Li lithium 7	4 Be beryllium 9											5 B boron 11	6 C carbon 12	7 N nitrogen 14	8 O oxygen 16	9 F fluorine 19	10 Ne neon 20		
11 Na sodium 23	12 Mg magnesium 24											13 Al aluminium 27	14 Si silicon 28	15 P phosphorus 31	16 S sulfur 32	17 Cl chlorine 35.5	18 Ar argon 40		
19 K potassium 39	20 Ca calcium 40	21 Sc scandium 45	22 Ti titanium 48	23 V vanadium 51	24 Cr chromium 52	25 Mn manganese 55	26 Fe iron 56	27 Co cobalt 59	28 Ni nickel 59	29 Cu copper 64	30 Zn zinc 65	31 Ga gallium 70	32 Ge germanium 73	33 As arsenic 75	34 Se selenium 79	35 Br bromine 80	36 Kr krypton 84		
37 Rb rubidium 85	38 Sr strontium 88	39 Y yttrium 89	40 Zr zirconium 91	41 Nb niobium 93	42 Mo molybdenum 96	43 Tc technetium 101	44 Ru ruthenium 101	45 Rh rhodium 103	46 Pd palladium 106	47 Ag silver 108	48 Cd cadmium 112	49 In indium 115	50 Sn tin 119	51 Sb antimony 122	52 Te tellurium 128	53 I iodine 127	54 Xe xenon 131		
55 Cs caesium 133	56 Ba barium 137	57–71 lanthanoids	72 Hf hafnium 178	73 Ta tantalum 181	74 W tungsten 184	75 Re rhenium 186	76 Os osmium 190	77 Ir iridium 192	78 Pt platinum 195	79 Au gold 197	80 Hg mercury 201	81 Tl thallium 204	82 Pb lead 207	83 Bi bismuth 209	84 Po polonium —	85 At astatine —	86 Rn radon —		
87 Fr francium —	88 Ra radium —	89–103 actinoids	104 Rf rutherfordium —	105 Db dubnium —	106 Sg seaborgium —	107 Bh bohrium —	108 Hs hassium —	109 Mt meitnerium —	110 Ds darmstadtium —	111 Rg roentgenium —	112 Cn copernicium —	113 Nh nihonium —	114 Fl flerovium —	115 Mc moscovium —	116 Lv livermorium —	117 Ts tennessine —	118 Og oganeson —		
lanthanoids																			
57 La lanthanum 139	58 Ce cerium 140	59 Pr praseodymium 141	60 Nd neodymium 144	61 Pm promethium —	62 Sm samarium 150	63 Eu europium 152	64 Gd gadolinium 157	65 Tb terbium 159	66 Dy dysprosium 163	67 Ho holmium 165	68 Er erbium 167	69 Tm thulium 169	70 Yb ytterbium 173	71 Lu lutetium 175					
actinoids																			
89 Ac actinium —	90 Th thorium 232	91 Pa protactinium 231	92 U uranium 238	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —					

SAFETY RULES

Personal protection

- Wear a laboratory coat and eye protection (goggles) all the time.
- Do not eat or drink inside the laboratory, not even chewing gum.
- Pipetting by mouth is strictly forbidden. Use the pipette fillers or bulbs that were supplied.
- Avoid laser beam to face your eye directly

Handling materials

- Avoid contact with reagents, you may wear the gloves provided for your skin protection.
- If you accidentally touch or spill any chemicals, wash yourself with running water as soon as you can.
- If any glass instrument breaks, do not handle the glass pieces yourself, ask the supervisor for help.
- If there is any other issue regarding your safety, please ask your supervisor to assist you.

Experiment



Q1-1

TWN-All (Chinese Taipei)

Determination of Refractive Index of Highly Concentrated Solutions of NaCl and Glycerin using Total Internal Reflection (TIR) (14 points)

Please read the general instructions in the separate envelope before you start this experiment.


Dependence of Refractive Index of a Solution on Concentration

Introduction:

Basic Principle: When a beam of light is incident on a surface, reflection and refraction take place. When the surface is uneven (rough), these phenomena result in scattering light in all directions. If the surface is that of a transparent medium, then most of the light is transmitted (refracted) and a small portion of it is reflected. If the surface is opaque and polished (like a metallic surface), all the light is reflected.

In this experiment, the green laser light is shone normally into the water inside a container. The container used has vertical walls. The laser beam first meets the smooth top water surface then it gets scattered from the rough bottom surface of container while producing a bright green spot on the bottom surface. This scattered light travels back into the water in all directions. This light again meets the top smooth surface of water and undergoes reflection, refraction and another phenomenon called total internal reflection (see Figure 1).

The scattered rays that reach the top surface of the water at an angle greater than the critical angle, get totally internally reflected which results in a bright ring enclosing a dark region.



(a) Air (b) Liquid (c) Laser Pointer (d) Container with Liquid

Figure1: (Left) Arrangement to observe the phenomenon. (Right) The ray diagram.

Experiment



Q1-2

TWN-All (Chinese Taipei)

From the definition of refractive index and critical angle we have:

$$\mu = \frac{1}{\sin(\theta_C)} = \frac{\sqrt{(\frac{d}{4})^2 + (h)^2}}{\frac{d}{4}} = \frac{\sqrt{(d)^2 + 16 \times (h)^2}}{d} \quad \text{--- (1)}$$

Where μ is the refractive index (RI) of liquid, d is the diameter of the dark disc and h is the height/depth of liquid. This formula can be applied to any transparent liquid medium.

From equation-(1)

$$(d)^2 \times (\mu)^2 = (d)^2 + 16 \times \frac{(V)^2}{(A)^2}$$

Where, A is the effective area of the horizontal cross-section of the container and V is the volume. $h = \frac{V}{A}$

The diameter (d) as a function of refractive index and area of the container (A) is

$$d = \frac{4}{A \times \sqrt{\mu^2 - 1}} \times V = S \times V \quad \text{--- (2)}$$

The diameter is proportional to volume and S is the proportionality constant given by

$$S = \frac{4}{A \times \sqrt{\mu^2 - 1}} \quad \text{--- (3)}$$

Using a liquid whose refractive index we know, we can calculate A . The effective area of the horizontal cross-section of the container is given by

$$A = \frac{4}{S \times \sqrt{\mu^2 - 1}} \quad \text{--- (4)}$$

The present experiment is to determine the refractive indices of salt and Glycerin solutions at certain high concentrations, using the refractive index of water (1.33).

Percent Concentration of Solutions:

Percent concentration **volume per volume** (V/V) is defined as the volume of solute in ml in 100 milliliters of solution. Hence 50% solution of any solute is 50 ml of solute in 100 ml of solution).

Experiment

Aim:

1. Determination of refractive index of 30% NaCl solution and Glycerin.
2. Determine the dependence of RI on the concentration of Glycerin water solution.

Equipments: You are supplied with the following equipment for this experiment:

Sr. No.	Item	Specifications	Quantity
01	Green LASER Pointer	Wavelength-532 nm	1 no + 1 spare
02	Burette stand	As shown in final assembly	1 no
03	Beaker	500 ml	3 no
04	Syringe	50 ml	1 no
05	Digital thermometer	To measure Room Temp	1 no
06	Glass stirrer	For making solutions	1 no
07	Container	As shown	1 no
08	Sodium Chloride (NaCl) Solution	30 % from AR grade salt	500 ml
09	Glycerin	AR grade	500 ml
10	Distilled water	Solution + washing	5000 ml
11	Tissue paper		
12	Safety goggle	Polaroid	1 no
13	Divider	Screw adjustable	1 no
14	Steel scale (ruler –optional)	0.5 mm Least Count	1 no
15	Reading lens	High quality	1 no

Warning :



Avoid direct eye exposure and through reflections.

Avoid staring at the laser spot for too long, advise to turn off the laser when it's not used for performing measurements

Experiment


The Experiment

Part-0 Measurement of the room temperature (0.2 points)


- A.0** Measure the room temperature using the thermometer provided and record (0.2pt) your reading in the answer sheet.
(Get the supervisor's sign after taking this reading)

Steps for setting up the equipment .

Step-1: Making a cardboard box.



(a) Cardboard Holder (b) Cardboard Box (c) Tape



Experiment


Wear your safety goggles all the time. If you are already wearing spectacles, wear the safety goggles above that. Do not look directly into the laser light.

Switch off the laser light when you are not taking the readings.

Glycerin should be kept covered when not in use.

Part 1. Calculation of effective area of cross section (A) of the container using distilled water (3.6 points)

Step-2: Attaching the Green LASER Pointer on the burette stand.
The laser pointer should be vertical .



(a) Cardboard Holder (b) Fisher Clamp (c) Laser (d) Opaque Rectangular container(TIR Container) (e) Stand Rod

Experiment

Step-3: Attaching the cardboard box on the burette stand.



(a) Cardboard Holder (f) Tape for support

Experiment



Q1-7

TWN-All (Chinese Taipei)

Procedure for Observation:

Apparatus set up:

Clamp the laser light in the burette stand in such a way that the switch is pressed by the middle grip of the stand. (When it is to be switched off, simply rotate the laser around the vertical axis so that the pressure on the switch is released. When it is to be switched on, rotate it back to the original position).

Make sure that the base of the stand on which the container is placed is horizontal.

Switch off the laser light when you are not taking reading.

Place the container below the light so that the laser beam falls on the bottom of the container.

(i) **Add 50 ml of distilled water** into the container. Switch on the laser, a bright ring with a dark disc inside becomes immediately visible.

- A.1** (ii) **Using the divider and ruler provided, measure the diameter of the dark disc.** Use the reading lens for better observation of the disc diameter. Record your readings in the **Table 1** of the answer sheet. (1.2pt)
(iii) **Repeat steps (i) and (ii)** by adding water in steps of 50 ml to **obtain six readings.**

- A.2** (iv) **Plot a graph on the given graph sheet (Graph 1 - plot 1)** with diameter of the dark disc (d) on the vertical axis and the volume (V) of water on the horizontal axis using the given graph page in the answer sheet. (1.8pt)
Note : (0,0) will be an additional data point to be plotted on the graph
(The total number of points to be plotted is 7)
while plotting use symbol
(.) dot = water
For marking the points on the graph

- A.3** (v) **Calculate the slope from the graph ($S = d/V$)** (0.2pt)

- A.4** (vi) **Calculate the effective area (A)** of cross-section of the container from the slope and equation (4) (0.4pt)

Experiment



Q1-8

TWN-All (Chinese Taipei)

Part 2: Determination of Refractive index of 30% NaCl Solution (3.4 points)

You are supplied 500ml of 30% NaCl Solution.

(i) Clean the container dry it by dabbing it with tissue paper.

B.1 (ii) Using the salt solution with fixed concentration, follow the steps (i) to (iii) of **Part 1**. Enter your readings in **Table 2** of your answer sheet. (1.2pt)

B.2 (iii) **Plot a graph on the same graph sheet** in the answer sheet (**Graph 1 - plot 2**) overlapping graph plotted in Part 1) with diameter on the vertical axis and volume on the horizontal axis. Label the points for distinction.
Note : (0,0) will be an additional data point to be plotted on the graph
(The total number of points to be plotted is 7)
while plotting use symbol
(+) plus = NaCl solution
For marking the points on the graph (1.6pt)

B.3 (iv) **Calculate the slope from the graph** (0.2pt)

B.4 (v) **Calculate the refractive index** of 30% NaCl solution from the slope and the value of **A** calculated in **Part 1**. (0.4pt)

Experiment



Q1-9

TWN-All (Chinese Taipei)

Part 3-A: Determination of refractive index of Glycerin (3.4 points)

You are supplied with 500 ml of Glycerin.

(i) Clean the container and dry it by dabbing it with tissue paper.

C-1.1 (ii) Using the pure glycerin provided **follow the steps (i) to (iii) of Part 1**. Enter your readings in **Table 3a** of your answer sheet. (1.2pt)

C-1.2 (iii) **Plot a graph on the same graph sheet** in the answer sheet (**Graph 1 - plot 3**) overlapping graph plotted in Part 1 and 2) with diameter on the vertical axis and volume on the horizontal axis. Label the points for distinction.
Note : (0,0) will be an additional data point to be plotted on the graph
(The total number of points to be plotted is 7)
while plotting use symbol
(*) star = Glycerin
For marking the points on the graph (1.6pt)

(iv) Do not disturb this solution at this point as it is required for part 3B of this experiment.

C-1.3 (v) Calculate the slope from the graph. (0.2pt)

C-1.4 (vi) Calculate the refractive index of glycerin from the slope calculated in this part of experiment and the value of A already calculated in Part 1 of this experiment. (0.4pt)

Experiment



Q1-10

TWN-All (Chinese Taipei)

Part 3B: Relation between Refractive index and concentration of Glycerin solution. (3.4 points)

Glycerin is miscible with water in all proportions; however, it takes thorough stirring to obtain a homogeneous mixture. In this part you will be measuring the refractive index of different concentrations of aqueous solutions of Glycerin.

(i) Using the syringe provided, remove 150 ml of Glycerin from the container, so the remaining amount of glycerin is 150 ml in the container.

- C-2.1** (ii) Measure d and enter values of volume and diameter in Table 3b in the answer sheet. (1.6pt)
(iii) Now add 50 ml of water to the container, stir the mixture gently and thoroughly to make a homogeneous solution.
(iv) Calculate the new concentration of the solution.
(v) Measure the diameter of the ring and record values of volume, diameter and concentration in Table 3b in the answer sheet.
(vi) Repeat steps (iii) to (v) for two more dilutions.

Calculate the values of S and Refractive Indices of the solutions and enter the values in the Table 3b in the answer sheet.

- C-2.2** (vii) **Plot the values refractive index on the vertical axis against the concentration on the horizontal axis (graph -2) in the answer sheet.** (1.4pt)
Note : (0,1.33) will be an additional data point to be plotted on the graph .
(The total number of points to be plotted is 5)

At this stage you have measured the refractive indices of 30 % NaCl solution and glycerin. You have also determined the relation between concentration and refractive index for glycerin solutions.

Answer the following questions in the answer sheet by choosing correct option:

- C-2.3** **How does the refractive index change with the concentration of glycerin solutions?** (0.2pt)
a. Increases with concentration
b. Decreases with concentration
c. Does not change with concentration

- C-2.4** **How would you expect the refractive index of NaCl solution to change with concentration?** (0.2pt)
a. Expected to increase with concentration
b. Expected to decreases with concentration
c. Expected not to change with concentration

Determination of the Glucose Content of Date Syrup Sample (8 Marks)

Please read the general instructions before you start this problem.


Date palms (*Phoenix dactylifera*) are found in abundance in the Middle East and in desert areas. The latin name of the tree is believed to have been derived from Greek *Phoenix daktulos*, which means purple or red finger. In the UAE, the late Sheikh Zayed Bin Sultan Al Nahyan was the founder of the country's agricultural renaissance. He encouraged the people of UAE to cultivate dates beyond the borders of the oasis where they traditionally grew dates, which played a great role in the flourishing dates throughout the nation. This great achievement turned the desert into an abundant paradise for its dwellers; UAE today brings this ancient super fruit to a new global market, which made the UAE the leader of date's cultivation in the world. From dates, a variety of food products are made.



Dates are rich in sugar which consists of two isomeric carbohydrates-Glucose (Molecular Formula $C_6H_{12}O_6$) and Fructose (Molecular Formula $C_6H_{12}O_6$). Glucose is the most important source of energy in all organisms.


Glucose contains an aldehyde group ($-CHO$) and hence is an aldose. Fructose contains a keto group ($-C=O$) and hence is a ketose. Glucose can be quantitatively oxidized to Gluconic acid ($C_6H_{12}O_7$) by iodine in alkaline medium. This enables the estimation of Glucose in presence by Fructose.

Date syrup is treated with iodine solution in the presence of Na_2CO_3 solution.



Glucose: Sodium hypoiodite = 1:1

The iodate (I) (hypoiodite) is converted back to iodine after adding acid.




On the completion of the oxidation, the excess iodine is back titrated against $Na_2S_2O_3$ solution using starch as an indicator.

Experiment



Q2-2

TWN-All (Chinese Taipei)



You are supplied with the following:

Chemicals	Labeled as	Quantity Supplied
Date Syrup	Date Syrup Sample	Supplied in Plastic Container
Iodine solution in a closed bottle	Iodine	100 mL
0.10 M $Na_2S_2O_3$ in a beaker	0.10 M $Na_2S_2O_3$	150 mL
15% Na_2CO_3 in a beaker	15% Na_2CO_3	50 mL
2M HCl in a beaker	2M HCl	100 mL
Starch	Starch Indicator	15 mL in Falcon Tube
Distilled Water	Distilled Water	1000 mL in a bottle
Apparatus	Labeled as	Quantity supplied
25 mL burette on stand	B ₁	1
10 mL volumetric pipettes	P ₁ , P ₂	2
Pipette fillers		2
250 mL Glass stoppered bottles		3
150 mL conical flasks	C ₁ to C ₃	3
Funnel		1
100 mL Volumetric flask	V ₁	1
10 mL measuring cylinder		1
Wash bottle with distilled water		1
Dropper		1
Beaker 150mL		1

- 10 mL measuring cylinder is to be used for 15% Sodium Carbonate, 2M HCl and starch indicator. Make sure that you wash it before every use.
- Feel free to use beakers, conical flasks and funnels that are available in your laboratory that fit with the experimental requirements, if necessary.

Procedure

i) Standardization of Iodine solution

Experiment



Q2-3

TWN-All (Chinese Taipei)

1. Rinse burette B_1 with 3 to 5 mL of 0.1 M $Na_2S_2O_3$
2. Fill burette B_1 with 0.1 M $Na_2S_2O_3$ using a funnel.

Note down the initial burette reading in Observation Table 1 in the answer sheet provided to you.

3. Using pipette P_1 take exactly 10 mL iodine solution in a 150 mL conical flask C_1 .
4. Using measuring cylinder add approximately 20 mL of distilled water to the Iodine solution taken.
5. Titrate the iodine solution against 0.1 M $Na_2S_2O_3$ from burette B_1 till a yellowish or light brown colour appears.
6. Add 2 mL of starch indicator using 10 mL measuring cylinder, the solution turns blue, and continue to titrate. The endpoint is colour change from blue to colourless.
7. Repeat the titration twice using conical flasks C_2 and C_3 .

2.1	Record your titration readings in Observation Table 1 and note down the burette reading.	(3.0pt)
------------	---	----------------

2.2	Calculate the molarity of Iodine solution.	(0.5pt)
------------	---	----------------

ii) Estimation of Glucose in Date Syrup


1. Using a funnel fill burette B_1 with 0.1 M $Na_2S_2O_3$.
2. **Note down the initial burette reading in Observation Table 2 in the answer sheet provided to you.**
3. Dissolve the date syrup supplied in the plastic container with warm distilled water. Transfer the solution quantitatively in the 100 mL volumetric flask V_1 and dilute with distilled water up to the mark using dropper. (You can use funnel to transfer the glucose solution.)
4. Insert the stopper and shake the solution to homogenise it. Transfer the homogenised solution to a correctly labelled 150 mL beaker.
5. Using pipette P_2 , take exactly 10 mL of the homogenised solution in a 250 mL glass stoppered bottle.
6. Using a 10 mL measuring cylinder, add approximately 10 mL of 15% Na_2CO_3 solution and add exactly 10 mL Iodine solution using pipette P_1 .
7. **Stopper the bottle and keep aside in the dark for about 30 minutes.** Repeat Steps 5 and 6 using two other 250 mL glass stopper bottles and keep them aside in the dark for about 30 mins each.
8. After 30 minutes, add 10 mL of 2M HCl using measuring cylinder, stopper the bottle and shake well (Iodine will liberate).
9. Add $Na_2S_2O_3$ drop wise from the burette B_1 till a light yellowish / light brown colour appears and then add 2 mL of starch solution indicator (dark blue colour will be obtained) using a 10 mL measuring cylinder and continue to titrate.
10. The endpoint is colour change from dark blue to colourless.
11. Repeat the titration twice.

Experiment



Q2-4

TWN-All (Chinese Taipei)



2.3 Record your titration readings in Observation Table 2 and note down the burette reading. (3.0pt)

Molar mass of Glucose=180 g/ mol

Ask for the mass of Date syrup given to you m(g)=..... g.

Observations

1. Calculate the number moles of Iodine solution added to 10 mL date syrup

=.....

2. Calculate the number of moles of Iodine which remain unused in the glucose oxidation reaction

=

.....moles

3. Calculate the number of moles of Iodine used up for oxidation of 10 mL date syrup

=.....moles

4. Calculate the total number of moles Iodine used up for the oxidation of the given date syrup

the number of moles of Iodine =.....moles

Experiment



Q2-5

TWN-All (Chinese Taipei)

2.4 Calculations

(1.5pt)

Determine a) the number of moles b) mass c) percentage of Glucose present in the given date syrup sample.

DELEGATION PRINT

Experiment



Q3-1

TWN-All (Chinese Taipei)

pH Titration using Universal Indicator (6 marks)

Please read the general instructions before you start this problem.

An acid base indicator is a substance that changes colour depending on the pH of the solution to which it is added. Such substances are used in solution or powder form for the determination of pH of a solution or to detect changes in pH during acid-base titrations.

A Universal indicator is able to change colours in a wide range of pH, and is hence used to determine how acidic or basic a solution is. Universal indicator that can be used to indicate the pH of the solution. It is a mixture of indicators that shows different colours at different pH values, and can be used in solution form or as paper strips. A colour chart supplied with the indicator solution or strips enables the determination of pH value.

pH range	Description	Colour
< 3	Strongly acidic	red
3 -6	Weakly acidic	orange, yellow
7	Neutral	green
8 -11	Weakly alkaline	blue
> 11	Strongly alkaline	violet

For the ease of the translations use the following color index to express the color as a letter, so it is easier for the markers to identify.

R	Red
O	Orange
Y	Yellow
G	Green
B	Blue
V	Violet

In case if you **ONLY** have Universal indicator as strips , use the following procedure. For every successive additions from the burette , stir the solution well, then touch the solution with a glass rod or dropper to take a very little of the solution and put it on the indicator strip and record the color. Repeat this until the end of the titration.

Warning: This procedure is written so that error in using pH strips is minimised. However, it will have an important error compared to the use of an indicator solution.

In this experiment, the equivalence point of an acid-base titration will be determined using Universal indicator to determine pH during the titration of weak acid with strong base.

You are supplied with the following:

Experiment

	Apparatus	Labelled as	Quantity supplied
1	10 mL pipettes	P_3, P_4, P_5 and P_6	4
2	25 mL burette	B_2	1
3	100 mL volumetric flasks	V_2, V_3, V_4	3
4	250 mL beaker		1
5	150 mL conical flasks	C_4, C_5	2
6	Funnel		1
7	Indicator bottles		2
8	Dropper*		1

* Use Dropper from Q-2

Chemicals	Labelled as	Quantity supplied
0.1M Succinic acid	0.1M Succinic acid	100 mL in a beaker
0.1M $NaOH$ (approx)	0.1M $NaOH$ (approx)	100 mL in a beaker
Weak acid solution	Weak acid	In 100 mL volumetric flask V_4
Universal indicator solution	Universal indicator with colour chart	In indicator bottle
Phenolphthalein indicator	Phenolphthalein	In indicator bottle
Distilled water	Distilled water	1000 mL in a bottle

- **Feel free to use beakers, conical flasks and funnels that are available in your laboratory that fit with the experimental requirements, if necessary.**

Procedure

1. Preparation of 100 mL of 0.01 M Succinic acid solution.

Using pipette P_3 take 10 mL of the supplied 0.1 M succinic acid solution in volumetric flask V_2 and dilute up to the mark using distilled water.

2. Standardisation of the diluted $NaOH$ solution.

1. Using pipette P_4 take 10 mL of the supplied 0.1 M (approx) $NaOH$ solution in volumetric flask V_3 and dilute up to the mark using distilled water.
2. Rinse burette B_2 with 3-5 mL of diluted $NaOH$.
3. Using a funnel fill the burette with diluted $NaOH$.

Note down the initial reading in Observation Table 1 in the answer sheet provided to you.

4. Using pipette P_5 take 10 mL of 0.01M Succinic acid solution in the conical flask C_4 and add 2 drops phenolphthalein indicator.
5. Titrate the solution against the diluted $NaOH$ till a faint pink colour persists.
6. Repeat the titration until you get three reasonable readings.

Experiment



Q3-3

TWN-All (Chinese Taipei)

3.1 Record your titration readings in Observation Table 1. Note down the reading. (1.5pt)

3.2 Molarity of $NaOH$ solution used in titration =M (0.5pt)

3. Titration of weak acid with strong base

1. Using a funnel fill burette B_2 with diluted $NaOH$
2. Dilute the weak acid solution supplied in volumetric flask V_4 to 100 mL using distilled water.
3. Using pipette P_6 take 10 mL of the diluted weak acid solution in a conical flask C_5 and add 4 drops Universal indicator.

In this titration, diluted $NaOH$ is added in instalments to the weak acid solution. After the addition of each instalment of diluted $NaOH$, record the colour of the solution, and pH from the chart supplied to you.

Put down pH value only when exact colour match is seen. If the colour is intermediate between colours in the chart, write down the pH range.

4. Add the burette solution in instalments of 0.5 mL each till the colour of the solution changes to purple. Thereafter take four more readings adding instalments of 0.5 mL each.

3.3 Record your observations in Observation Table 2 in the answer sheet provided to you. (2.5pt)

3.4 Plot a graph of pH vs volume of diluted $NaOH$ and determine the 5 mL range of the equivalence point. (0.5pt)

3.5 Find ΔpH for every successive change in volume (0.5mL) and then plot a graph of $\Delta pH/\Delta V$ vs volume of diluted $NaOH$ only in the range identified in 3.4 above (0.5pt)

3.6 Determine the equivalence point from the data above. (0.5pt)

Experiment




Q4-1


TWN-All (Chinese Taipei)

General Instructions :

1. Wherever asked to mark the cell with a cross (X), mark as follows.



2. Wherever asked to mark the cell with a dash (-), mark as follows.



Q.4. Investigating mixup of newborn babies in a hospital

This is an investigative experiment to identify parents of three newborn babies who got mixed in a hospital. The identification will be done with the help of blood groups. Until the 1980s blood groups were used in forensics, but this has now been replaced by other techniques that are more reliable. ABO blood typing is still important for blood transfusions.

The presence or absence of A and B antigens on the Red Blood Cells (RBCs) determines the ABO blood group of an individual. A person with only antigen A will be of blood group A. A person with only antigen B will be of blood group B. A person with both antigens A and B will have blood group AB, while one who has neither of the antigens has a blood group of O type. These antigens in an individual are governed by the alleles of the gene responsible for their synthesis. Antigen A is determined by the allele I^A , while antigen B is determined by the allele I^B . No antigens are produced when an individual carries the i allele. The alleles I^A and I^B are co-dominant. The allele i is recessive to both I^A and I^B alleles.

The presence of a given antigen can be identified by the use of antibodies. For example, if an antibody against antigen A is added to blood from a person with blood group A, RBCs will clump together or agglutinate. In this experiment, you will not use blood samples but solutions that mimic the process of agglutination (clumping) of blood in the presence of a given antibody. The process has been mimicked using chemicals and precipitation represents the process of agglutination. The use of chemicals to demonstrate the concept of blood groups was developed by Magdalena Wajrak of Edith Cowan University in Perth, Australia (Harrison T, 2015, Science in school, 32: 33-36)

Materials provided:

Glassware and miscellaneous items

1. Mimicked blood samples (13 in all) in 1.5 ml plastic tubes (in stand) labelled as follows:
 - (i). Four samples W, X, Y and Z (to be used in Exercise 1).
 - (ii). Nine samples C, D, E, 1F, 1M, 2F, 2M, 3F, 3M (to be used in Exercise 2).
2. 3 X 15 ml plastic tubes labeled as Anti -A, Anti - B and NA which have mimicked antibodies against antigen A, antigen B and no antibody, respectively. These have been placed in a 250 ml plastic beaker.
3. Four Cavity slides with 3 wells in each.
4. 7 plastic droppers.
5. 250 ml beaker with distilled water

Experiment

6. Permanent marker
7. Rectangular labels
8. A4 sheet of black paper for placing slides
9. Waste bin
10. Additional distilled water in a bottle

Note: Please raise your hand, if you require additional distilled water.

Tissue papers and waste beaker will be provided by the supervisor

Exercise 1: Identify the blood groups of the blood samples W, X, Y and Z

1. Arrange four cavity slides to make a grid similar to the one shown in Figure 1.1. The slides should be placed on the black sheet provided to you.




Figure 1.1

2. With the marker pen and labels provided, mark the wells of the grid as above. NA stands for no anti-body.

3.1 Mark three droppers one as Anti-A (to be used only for taking Antibody A), one as Anti-B (to be used only for taking Antibody B) and one as NA (to be used for taking the solution marked as NA).

3.2 Use the remaining droppers for blood samples.

3.3 Before starting and in between taking two different samples, flush the dropper several times (15 to 20 times) with distilled water to ensure that they are clean.

3.4 Ensure that the droppers are clean so that samples do not become cross-contaminated. Please avoid touching the added solutions in any of the wells of the cavity slide.

4. Using a clean dropper, place a drop of blood sample W in each of the wells in column 1.
5. Continue the same with the other three blood samples (X, Y and Z).
6. In the first row, add 1 drop of Anti A (antibodies against A-antigen) in each of the wells.
7. In the second row, add 1 drop of Anti B (antibodies against B-antigen) in each of the wells.
8. In the third row, add 1 drop of NA solution.

Experiment

- A.1.1** Observe the wells and record the presence of white precipitate (mimicking agglutination of blood) in the **appropriate cells in Table 1.1 by marking a cross (X)**. Mark a dash (-) in cells representing wells where no precipitation was observed.

Table 1.1				
	W	X	Y	Z
Anti-A				
Anti-B				
NA				

- A.1.2** Request your supervisor to take a photograph of the plate. The supervisor will submit the photograph on your behalf. (0.25pt)

Based on your observation, identify the blood groups of the samples W, X, Y and Z in Table 1.2 by **marking a cross (X) in the appropriate cell**.

A.1.3

(0.25pt)

Table 1.2				
	Blood Group			
Sample	A	B	AB	O
W				
X				
Y				
Z				

Identify the row(s) (Anti A, Anti B and NA) in Figure 1.1 that act(s) as the control for the experiment with a **cross (X) in the correct cell(s)**. Mark dash (-) in the remaining one(s).

Experiment

A.1.4

(0.25pt)

Anti-A	Anti-B	NA

Exercise 2: Identify the blood groups of the parents and the babies in an attempt to restore the babies to their respective parents.

There are three newborn babies (C, D and E), whose tags indicating their parents have been mixed up. In order to identify the correct parents of the three babies, blood samples were taken from the babies and the possible parents (1 to 3). The experiment attempts to identify the parents of the three babies based on their blood groups.

For the identification you are provided with mimic of 9 blood samples labeled as follows:

Note: In the table, F stands for Father (not female) and M stands for Mother (not male).

Sample No.	Label	Blood sample from
1	1F	Parent 1 father
2	1M	Parent 1 mother
3	2F	Parent 2 father
4	2M	Parent 2 mother
5	3F	Parent 3 father
6	3M	Parent 3 mother
7	C	Baby C
8	D	Baby D
9	E	Baby E

2.1. For each of the blood samples, identify the blood group following the procedure described in the first exercise.

Note:

- Before reusing the cavity slides, wash them carefully with distilled water and dry them well with tissue paper before putting samples in them.
- Ensure that the droppers are clean before taking the blood samples.

In Table 1.3, **mark a cross (X) in the appropriate cells** for the presence of precipitate. Mark a dash (-) in cells representing wells where no precipitation was observed.

Experiment

A.2.1.1 Take photographs of the labeled slides as done in 1.2. Request your supervisor to take photographs of the labeled slide. The supervisor will submit the photograph on your behalf. (4.5pt)

Table 1.3									
	1F	1M	2F	2M	3F	3M	C	D	E
Anti-A									
Anti-B									
NA									

Based on the results of the experiment, identify the blood groups of each of the nine samples by **marking a cross (X) in the appropriate cell** in Table 1.4.

A.2.1.2

(0.50pt)

Table 1.4				
Blood group of babies				
Baby	Blood group A	Blood group B	Blood group AB	Blood group O
C				
D				
E				
Blood group of parents				
1F				
1M				
2F				
2M				
3F				
3M				

Based on the different blood groups identified by you, indicate the possible parent pairs for the babies C, D and E by marking a cross (X) in the appropriate cell(s) in Table 1.5. There could be more than one possibility. Mark a dash (-) in the remaining cell(s).

Experiment

A.2.2

(1.0pt)

Table 1.5			
	Parent pair 1	Parent pair 2	Parent pair 3
Baby C			
Baby D			
Baby E			

Based on your interpretations of the blood groups which baby(ies) (C, D, E) can be matched to their parent pairs (1, 2, 3) with certainty based on the evidence?

Write the number of the parent pair (1, 2 or 3) in the cell to the corresponding baby(ies).

Mark a dash (-) in cell(s) corresponding to a child with multiple possible parent pairs.

A.2.3

(0.25pt)

Child	C	Parent	
Child	D	Parent	
Child	E	Parent	

Predict the genotype of the child and the parent pair that can be matched with certainty based on your answer in 2.3.

Indicate **one possible genotype** of the child and the corresponding parent pair.

Mark a dash (-) in cell(s) corresponding to a child with multiple possible parent pair.

A.2.4

(0.25pt)

		Genotype of the child			Genotype of the parents	
					Father	Mother
Child	C		Parent			
Child	D		Parent			
Child	E		Parent			

Q.5. Analyzing human chromosomes

The karyotype of a species represents the chromosomes of a cell, usually displayed as a systematic arrangement of chromosome pairs in descending order of size. In order to make a human karyotype, metaphase chromosomes are prepared from blood cells

The blood cells are induced to divide in culture and then treated with colchicine to block cell division at metaphase. The cells are then broken by hypotonic treatment and chromosomes are spread on glass slides. The chromosomes are stained and observed under a microscope. The photograph of the metaphase (as shown in Figure 5.1) is then used to make a karyotype. Karyotypes can be used to identify chromosomal abnormalities and aberrations. In the manual method, individual chromosomes are cut from the photograph and then lined up by size and the position of the centromere with their respective partners. In humans chromosomes can be of three types, which are determined by the position of the centromere (point of attachment of the mitotic spindle): (i) **acrocentric chromosomes** have the centromere located very close to the end resulting into one of the arms being very short (sometimes not even discernable), (ii) **submetacentric chromosomes** have arms of unequal lengths and (iii) **metacentric chromosomes** have equal or almost equal arms. The karyotype prepared from the metaphase spread shown in Figure 5.1 is presented in Figure 5.2. A description of the chromosomes belonging to different groups (Figure 5.3) is given in Table 5.1.



Figure 5.1. A metaphase spread of human chromosomes

Experiment




Figure 5.2. Karyotype prepared from the metaphase spread presented in Figure 5.1.

Group	Chromosomal Pairs	Description
A	1-3	Large almost metacentric chromosomes
B	4-5	Large submetacentric chromosomes
C	6-12 + X	Medium-sized submetacentric chromosomes
D	13-15	Large acrocentric chromosomes
E	16-18	Small submetacentric chromosomes
F	19-20	Small metacentric chromosomes
G	21-22 + Y	Small acrocentric chromosomes
XY	X	Medium-sized sub metacentric chromosome
	Y	Small acrocentric chromosome

In the following exercise you are required to prepare a karyotype from the photograph of the metaphase spread given to you. This chromosomal preparation is from an individual with an anomaly in their sex chromosomes.

Experiment



Q5-3

TWN-All (Chinese Taipei)

Materials given for karyotype preparation:

1. Photograph of a metaphase spread for karyotype preparation
2. Plastic petri-dish
3. Fine scissors
4. Forceps
5. Transparent tape
6. Sheet marked 'Karyotype' to paste the chromosome cutouts

Procedure:

Use the photograph of the metaphase spread for the following exercise

Exercise 1: Count the number of chromosomes

A.5.1 Count the number of chromosomes and record in the answer book.

(0.25pt)

Experiment

Exercise 2: Make a karyotype

1. With the help of fine scissors cut out each chromosome and place it in the given petri- dish. Make sure that you do not loose any of them.
2. Arrange the chromosomes (cut outs) on the sheet labeled **karyotype** according to the size of the chromosomes and position of their centromere. Use Figure 5.3 and Table 5.1 as your reference. After arranging the chromosomes stick them in place with transparent tape. Take a photograph and attach to the answer sheet.
3. **The sheet labeled karyotype is part of your answer book and is placed at the end of the answer book.**

In each group, arrange the chromosomes by their approximate length. There will be no penalty for errors in identification of specific chromosomes within a group.

Figure 5.3 is a pictorial guide on how to prepare a karyotype.




Figure 5.3 A pictorial representation of preparing a karyotype. 1: Items to be used for making a karyotype, 2-5: Cutting the individual chromosomes, 6-8: Picking the chromosomes with the fine forceps and collecting them, 9: Arranging the chromosomes, 10-11: Sticking the chromosomes with the piece of tape.

A.5.2 Please ask your supervisor to either scan or take a photograph of the karyotype (3pt) on the answer sheet and submit.

Experiment

Exercise 3: Answer the following.

Can the following cells in human blood be used for preparation of metaphase chromosomes?

A.5.3.1 Mark a cross (X) in the appropriate column (Yes / No).

(0.25pt)

S.No.	Cells	Yes	No
1.	Erythrocyte (Red Blood Cells)		
2.	Lymphocyte (White Blood Cells)		

You are asked to prepare chromosomal spread from mitotic plant cells. Can you use the following plant parts to successfully prepare the chromosomal spread?

A.5.3.2 Mark a cross (X) in the appropriate cells (Yes/No).

(0.25pt)

S.No.	Plant parts	Yes	No
1.	Leaf blade		
2.	Anther		
3.	Root tip		

The photograph represents chromosomes in a rodent cell undergoing division. All chromosomes in this rodent are **acrocentric**.

Experiment

Q5-6

TWN-All (Chinese Taipei)



Does the following photograph represent the following stages of division?

A.5.3.3 Mark a cross (X) in the appropriate column (Yes/No).

(0.25pt)

Stage of division	Yes	No
Mitotic Metaphase	☐	☐
Mitotic Anaphase	☐	☐
Meiotic Metaphase I	☐	☐
Meiotic Anaphase I	☐	☐
Meiotic Metaphase II	☐	☐
Meiotic Anaphase II	☐	☐